

WETLAND MITIGATION SITE MONITORING REPORT
Harrison Avenue Extension at Pecatonica River Forest Preserve
Winnebago County, 2006

INTRODUCTION

This report details monitoring of the wetland mitigation site created to compensate for wetlands that were impacted during an extension of Harrison Avenue in Rockford, Illinois. Impacted wetlands totaled 1.82 ha (4.49 ac) in area, and included 0.66 ha (1.64 ac) of palustrine emergent wetland, 1.05 ha (2.60 ac) of forested wetland, and 0.10 ha (0.25 ac) of streambed. The compensation site is owned by the Winnebago County Forest Preserve District, and is located just south of Blair Road, adjacent to the Pecatonica River (legal location NW/4, Sect. 19, T 27N, R 10E). The site is approximately 24.3 ha (60 ac) in size, and was formerly agricultural land and a drainage ditch. Because the compensation site is located more than one mile from the impact site, a total of 7.27 ha (17.96 acres; a 4:1 ratio of compensated to impacted area) is required under the Illinois Interagency Wetland Policy Act of 1989. A total of 4.22 ha (10.44 ac) wet prairie and emergent marsh, 2.64 ha (6.52 ac) of forested wetland, and 0.40 ha (1.00 ac) of streambed wetland are planned for the site. The forested wetland area is located in southernmost part of the site, adjacent to existing forest along the Pecatonica River. The wet prairie and marsh are located in northern part of site near Blair Road. Wetland vegetation was also planted in an existing tributary along the west side of the property. Herbaceous wetlands were planted with seeds and live plant material, and the forested site was planted with wet prairie seeds and tree saplings (Hey and Associates 1999). Wetland construction was completed in 2003. On-site monitoring was conducted on 10, 17 and 24 August 2005 and 14 August 2006.

This report discusses the goals, objectives, and performance criteria for the mitigation project, the methods used for monitoring the site, the monitoring results from August 2006, and a discussion and recommendations based on those results. Methods and results are discussed by performance criteria for each goal.

Goals, Objectives, and Performance Standards

Goals, objectives, and performance standards follow those specified in the wetland compensation plan that Hey and Associates, Inc. (1999) listed for this site. Each goal should be attained by the end of the 5-year monitoring period. Goals, objectives, and performance criteria are listed below.

Project goal 1: Each wetland community should be a jurisdictional wetland as defined by current federal standards.

Objective: 4.22 ha (10.44 ac) of wet prairie and emergent marsh, 2.64 ha (6.52 ac) of forested wetland, and 0.40 ha (1.00 ac) of streambed wetland will compensate for impacts

to 0.66 ha (1.64 ac) of palustrine emergent wetland, 1.05 ha (2.60 ac) of forested wetland and 0.10 ha (0.25 ac) of streambed, at a total ratio 4 ha of wetland compensation to 1 ha of impacted wetland.

Performance criteria:

- a. Predominance of hydrophytic vegetation: More than 50% of the dominant plant species must be hydrophytic.
- b. Presence of wetland hydrology: The area must be either permanently or periodically inundated at average depths less than 2 m (6.6 ft) or have soils that are saturated to the surface for at least 12.5% of the growing season.
- c. Occurrence of hydric soils: Hydric soil characteristics should be present, or conditions favorable for hydric soil formation should persist at the site.

Project goal 2: The wetland plant community should meet minimum standards for floristic integrity and plant species composition.

Objectives: An emergent marsh, wet prairie, forested wetland, and a vegetated streambed wetland will be established by planting native wetland vegetation.

Performance criteria:

- a. Floristic Quality Index: Floristic Quality Index (FQI) (Taft et al. 1997) will be greater than or equal to 20 over entire project area by fifth year of site monitoring. Native FQI will increase each successive year after planting through the end of monitoring.
- b. Vegetation cover and species richness: By end of the fifth year the site will exhibit minimum of 60% vegetative cover and will include minimum of 30 native species.
- c. Mean wetness coefficient: Mean native wetness coefficient (Reed 1988) will be less than or equal to zero by end of the fifth year.
- d. Importance of native plant species: Relative importance value of total native plants will increase each successive year after planting.
- e. Non-native and weedy species dominance: By end of fifth year no more than 20% of the area will have non-native or weedy species dominance (including *Typha* spp., *Salix interior*, and *Phragmites australis*).

Project goal 3: The forested wetland should meet minimum standards for planted tree and shrub survival.

Objectives: A forested wetland will be established by planting native trees and shrubs.

Performance criteria:

- a. Percent survival of planted trees and shrubs: Planted tree and shrub survival must be at least 80% each year.
- b. Number of planted tree and shrub species: The site must have at least five species of planted trees and three species of planted shrubs.

METHODS

Project goal 1

a. Predominance of hydrophytic vegetation

The method for determining dominant vegetation at a wetland site is described in the *Corps of Engineers Wetlands Delineation Manual* (Environmental Laboratory 1987) and further explained in the *Federal Manual for Identifying and Delineating Jurisdictional Wetlands* (Federal Interagency Committee for Wetland Delineation 1989). The relative Importance Value, a combination of relative coverage and relative frequency, for each species was determined by quantitatively sampling vegetation at each site (see project goal 2, below). Species were then arranged by Importance Value in decreasing order, and Importance Values were sequentially summed, starting with the most prevalent species, until the total reached 50. Those species included in the summation were considered dominant species. Each of the dominant plant species was then assigned its wetland indicator status rating (Reed 1988). Any plant rated facultative or wetter (*i.e.*, FAC, FAC+, FACW, or OBL) is considered a hydrophyte. A predominance of vegetation in the wetland plant community exists if more than 50% of the dominant species present are hydrophytic.

b. Presence of wetland hydrology

In April 2005, Illinois State Geological Survey (ISGS) personnel installed a hydrologic monitoring network at the site (Plankell and Benton 2006). Locations of monitoring instruments at the site can be found in the ISGS report *Pecatonica River Forest Preserve Wetland Compensation Site* (Plankell and Benton 2006). Methods are further described in the ISGS document *Annual report for active IDOT wetland compensation and hydrologic monitoring sites* (Fucciolo et al. 2006).

c. Occurrence of hydric soils

The soil was sampled in each plant community in order to monitor hydric soil development. Soil profile morphology, including horizon color, texture, and structure, was described from one or more representative, permanent sampling points in each vegetation community. Additionally, the presence, type, size, and abundance of redoximorphic features were noted.

Hydric soils typically develop slowly, and characteristics may not be apparent during the first several years after project construction. In the absence of hydric soil indicators at the end of the five-year monitoring period, hydrologic data could be used as corroborative evidence that conditions favorable for hydric soil formation persist at the site.

Project goal 2

Vegetation in the wet prairie/emergent marsh community (hereafter referred to as Site 1) was quantitatively sampled using 0.5-m x 0.5-m (0.25 m²) quadrats (n = 29) placed every 30 m along five transects. A baseline was established along the northern boundary of the site, parallel to Blair Road. Five parallel transects placed every 30 m (100 ft) and running north to south were used to sample the community. Locations of these transects are marked on the enclosed map. All plant species in each quadrat were recorded and each species was assigned a cover class (Table 1), an estimate of the amount of area within the sample quadrat that is covered by that species. Data from quadrats were used to calculate frequency (percent of quadrats in which the species is present), relative frequency (frequency relative to other

species), average cover per quadrat, relative cover, and Importance Value (average of relative frequency and relative cover) for each sampled species.

Table 1: Cover classes used to estimate aerial cover by plant species in sample quadrats

Cover class	Range of aerial cover	Midpoint of range
R	<1%, solitary	0%
+	<1%, seldom	0%
1	1-5%	3%
2	5-25%	15%
3	25-50%	37.5%
4	50-75%	62.5%
5	75-95%	85%
6	95-100%	97.5%

Because the forested wetland community (Site 2) has been unsuccessful in meeting goals for wetland establishment and planted tree survival, herbaceous vegetation at site 2 was not quantitatively sampled in 2005 or 2006.

In order to approximate relative species abundances in the streambed wetland (Site 3), we employed a semi-quantitative method in which all species were assigned an abundance value ranging from one to five based on visual estimates of abundance and cover. Species that were present as a single or a few individuals were assigned a value of one; species present, but uncommon were assigned a two; species which occurred at moderate frequency throughout the site were assigned a three; species which were common and abundant across the site were assigned a four; and species which were dominant at the site (greater than 20% areal cover) were assigned a five.

The Floristic Quality Assessment (Taft et al. 1997) was applied to each plant community at each site to evaluate ecological integrity. The assessment methodology is used to identify natural areas and facilitate floristic comparisons among sites. This technique is part of the procedure for the long-term monitoring of natural areas and the monitoring of restored or created wetlands (Swink and Wilhelm 1994). Plant species not native to Illinois are not included in the FQI. Each native plant species is assigned a coefficient of conservatism (*C*) ranging from 0 to 10. Lower numbers have been assigned to species that tend to be more tolerant of disturbance and higher numbers to species that are generally found in less disturbed natural areas. A mean coefficient value (*mCv*) is determined by summing the coefficients of conservatism (*C*) and dividing by the total number of native species (*N*). The Floristic Quality Index (FQI) is then determined by dividing the sum of the coefficients of conservatism by the square root of *N*. This calculation is done to incorporate numerical species diversity into the FQI value. Sites with FQI values less than 10 suggest that the area has been highly disturbed or is in an early successional stage. Sites with FQI values of 20 or more generally possess some evidence of natural character and may be considered environmental assets. Sites with values of 35 or more are considered to be of natural area quality.

At each plant community the mean wetness coefficient was calculated as the average of the numerical values associated with individual species wetness coefficients for all species in the community (Reed 1988), where the numerical values are: UPL = 5, FACU- = 4, FACU = 3, FACU+ = 2, FAC- = 1, FAC = 0, FAC+ = -1, FACW- = -2, FACW = -3, FACW+ = -4, and OBL = -5.

Boundaries of the plant communities within the project area were recorded using a Trimble Global Positioning System. Locations of these sites were overlain on digital ortho quads (DOQs), and approximate wetland acreage was determined for each site using ArcView 3.2. Printouts of these DOQs are included with this report.

Project goal 3

All living planted trees and shrubs were counted and identified to species to assess survival rates.

RESULTS

Project goal 1

a. Predominance of hydrophytic vegetation

An emergent marsh/wet prairie community was planned for the northwestern portion of the project area (Site 1). During the 2006 monitoring season, a portion of Site 1 supported wet meadow and marsh vegetation, and is referred to as Site 1A. The remainder of Site 1, however, lacked dominant hydrophytic vegetation, and is referred to as Site 1B (see Fig. 2, Appendix B). The plant community at Site 1B is currently non-native grassland. Dominant plant species for the marsh/wet meadow (Site 1A), the non-native grassland (Site 1B), the forested site (Site 2) and the streambed wetland (Site 3) are shown in Tables 2, 3, 4 and 5. At two of the four sites (Sites 1A and 3), greater than 50% of the dominant species are rated OBL, FACW or FAC, and therefore, the dominant vegetation is hydrophytic.

Table 2. Dominant plant species by stratum for the marsh/wet meadow portion of Site 1A

Dominant plant species	Stratum	Indicator status
1. <i>Phalaris arundinacea</i>	herb	FACW+

Table 3. Dominant plant species by stratum for the non-native grassland portion of Site 1B

Dominant plant species	Stratum	Indicator status
1. <i>Chenopodium album</i>	herb	FAC-
2. <i>Phalaris arundinacea</i>	herb	FACW+
3. <i>Polygonum pensylvanicum</i>	herb	FACW+
4. <i>Setaria faberi</i>	herb	FACU+

Table 4. Dominant plant species by stratum and wetland indicator status for the forested site (Site 2)

Dominant plant species	Stratum	Indicator status
1. <i>Panicum virgatum</i>	herb	FAC+
2. <i>Polygonum pensylvanicum</i>	herb	FACW+
3. <i>Setaria faberi</i>	herb	FACU+
4. <i>Solidago canadensis</i>	herb	FACU

Table 5. Dominant plant species by stratum and wetland indicator status for the streambed wetland (Site 3)

Dominant plant species	Stratum	Indicator status
1. <i>Bidens frondosa</i>	herb	FACW
2. <i>Leersia oryzoides</i>	herb	OBL
3. <i>Lemna minor</i>	herb	OBL

b. Presence of wetland hydrology

Hydrologic data for the sites for September 2005 through September 2006 are presented in Appendix B. An estimated 2.5 ha (6.2 ac) within the total project area conclusively satisfied the wetland hydrology criterion for greater than 12.5% of the growing season during the monitoring period (Fig. 1, Appendix B; Plankell and Benton 2006). All of the area that satisfied the wetland hydrology criterion was located within the northern part of the project area (Site 1). Site 3, the streambed wetland, was not monitored by the ISGS.

c. Occurrence of hydric soils

Soils in the northwestern portion of the project area (Site 1) were mapped as four different soil map units (107, 242, 415, and 451) by the Natural Resources Conservation Service (NRCS) (Grantham 1980). Map unit 107 (Sawmill silty clay loam) was found in the western, wetter part of the site (Site 1A), whereas the southeastern, drier part of the site (Site 1B) was comprised of the map unit 415 (Orion silt loam). Map units 242 (Kendall silt loam) and 451 (Lawson silt loam) were not found during field investigation.

Sawmill silty clay loam is a poorly drained, hydric soil, formed in silty and clayey water-deposited sediments. It is typically found on low stream terraces, broad alluvial valleys, and small upland drainageways. Orion silt loam is a somewhat poorly drained, non-hydric soil. It formed in silty alluvium and is typically found along small streams and at the upper end of drainageways.

The NRCS mapped units 107 (Sawmill silty clay loam) and 451 (Lawson silt loam) in southern part of the project area (Site 2). Only map unit 451 was found during field investigation. Lawson silt loam is a somewhat poorly drained, non-hydric soil formed in alluvium. It is typically found on bottom lands along the Pecatonica River and along small streams and drainage ways.

Soil within the streambed wetland (Site 3) showed features of a hydric soil. The areas adjacent to the stream were mapped as Sawmill silty clay loam (map unit 107). We classified the soil of the stream bottom as a Vertic Epiaquept. The soil on 13 August 2006 was saturated in the top 8 cm (3 in) and ponded in a few places along the creek bed.

Typical pedons found within the project area are described below. The locations of the soil profiles used for the descriptions are marked on the enclosed map.

Table 6. Soil at the marsh/wet meadow (Site 1A) (Sawmill silty clay loam, hydric)

Depth [cm]	Matrix Color	Redox Concentrations	Redox Depletions	Texture	Structure
0 – 13	10YR 3/1	7.5YR 3/4	-	silty clay loam	subangular blocky
13 – 60	10YR 2/1	-	10YR 5/1 & 10YR 4/2	silty clay loam	subangular blocky

Table 7. Soil at the non-native grassland (Site 1B) (Orion silt loam, non-hydric)

Depth [cm]	Matrix Color	Redox Concentrations	Redox Depletions	Texture	Structure
0 – 33	10YR 3/2	-	-	silt loam	subangular blocky
33 – 60	10YR 2/1 & 10YR 4/2	10YR 3/6	-	silt loam	subangular blocky
60 – 81	10YR 3/1	10YR 3/6	10YR 4/2	silt loam	subangular blocky
81 – 91	10YR 3/1.5	7.5YR 3/4	-	silt loam	weak platy

Table 8. Soil at the forested site (Site 2) (Lawson silt loam, non-hydric)

Depth [cm]	Matrix Color	Redox Concentrations	Redox Depletions	Texture	Structure
0 – 33	10YR 2/1	-	-	silt loam to loam	angular blocky
33 - 119	10YR 3/2	-	-	silt loam to loam	subangular blocky to massive
119 - 130	Multi-colored	-	-	sand	single grain

Table 9. Soil at the streambed wetland (Site 2) (generic Vertic Epiaquept, hydric)

Depth [cm]	Matrix Color	Redox Concentrations	Redox Depletions	Texture	Structure
0 – 8	2.5Y 3/1	7.5YR 3/4	-	silt loam	massive
8 – 46	2.5Y 2.5/1	10YR 3/6	10YR 4/2	silt loam	subangular blocky
46 – 61	10YR 2/1	10YR 3/4	-	silt loam	massive

Project goal 2

Mean coefficient of conservatism and FQI values were calculated for each site from the species lists included in Appendix A. For each site, mCv and FQI values were calculated using only species that became established on the site naturally (volunteer species), and then recalculated to include planted species (Table 10). When planted species are included, the FQI currently exceeds the stated performance criterion of 20 for the entire project area. However, because the entire project area is large and includes several plant communities, it is likely to support a large number of native plant species regardless of the area's floristic conservation value. An FQI value above 20 for the entire project area should not be considered exceptionally high. When considered individually, FQI and mCv values for each

site suggest fair natural quality when planted species are included. As required by the performance criterion, native FQI increased from 2005 to 2006 for the entire project area.

Table 10. Mean coefficient of conservatism (mCv) and Floristic Quality Index (FQI) values for constructed wetlands

<u>a. Planted species not included</u>				
Site	2005		2006	
	mCv	FQI	mCv	FQI
1A. Wet meadow	2.3	12.9	1.9	13.0
1B. Non-native grassland	1.4	7.8	1.5	10.9
2. Forested site	1.8	11.0	1.7	10.8
3. Stream channel	2.1	11.3	2.5	13.5
TOTAL PROJECT AREA	2.1	17.9	2.0	18.9

<u>b. Planted species included</u>				
Site	2005		2006	
	mCv	FQI	mCv	FQI
1A. Wet meadow	2.8	17.7	2.4	17.7
1B. Non-native grassland	2.1	13.7	1.9	14.1
2. Forested site	2.6	19.6	2.6	19.6
3. Stream channel	2.4	13.8	2.7	15.8
TOTAL PROJECT AREA	2.7	27.1	2.7	28.4

All sites met the performance criterion of at least 60% vegetative cover in 2006. Areal cover by bare ground was less than 5%, on average, per 0.25 m² quadrat in Sites 1A and 1B. Though not quantified in Site 2, a similar amount of bare ground was observed in this site. The streambed wetland had large areas of bare ground, as to be expected from a streambed, but did have at least 60% vegetation coverage.

When planted species are included, a total of 116 native plant species were observed across the entire project area, far exceeding the performance criterion of 30 native species. However, the stated performance criterion is far too low for an area of this size, and is therefore not an appropriate standard for judging the floristic integrity of the area. When considered individually, each site had at least 30 native species (Site 1A: 46 species; Site 1B: 50 species; Site 2: 41 species; Site 3: 30 species).

The project area as a whole, and each site individually, met the performance criterion of having a mean wetness coefficient less than zero (total project area: -1.30; Site 1A: -2.19; Site 1B: -0.25; Site 2: -0.25; Site 3: -3.54).

The total Importance Values of native species was 40.0 at Site 1A and 32.1 at Site 1B, indicating a large amount of cover by, and high frequency of, exotic species. Two exotic grasses were particularly common: *Phalaris arundinacea* had an Importance Value of 52.3 at Site 1A, and *Setaria faberi* had an Importance Value of 33.6 at Site 1B.

The only dominant species in Site 1A, *Phalaris arundinacea*, is exotic. Three of the four dominant species at Site 1B, *Setaria faberi*, *Chenopodium album* and *Phalaris arundinacea*, are exotic. *Setaria faberi* is also a dominant at Site 2. All dominant species in Site 3 are native, non-weedy species. Therefore, only Site 3 met the performance criterion of having no more than 20% of the area with non-native or weedy species dominance.

Project goal 3

Seven species of planted native trees and five species of planted native shrubs were represented by live individuals at Site 2 in August 2006 (Table 11). Therefore, the site met the performance standard of having at least five species of planted trees and three species of planted shrubs. However, total tree and shrub survival was only 21.9%, far below the specified performance standard of 80% survival. A large number of *Juglans nigra* and *Fraxinus pennsylvanica* individuals have naturally regenerated at the site, making it difficult to distinguish naturally recruited versus planted individuals. The reported values for survival of planted individuals for these species are therefore overestimates. In addition, a large number of naturally recruited *Ulmus americana* seedlings were observed.

Table 11. Planted tree and shrub survival at Site 2

Species	Number planted	Surviving 2005	Surviving 2006	Percent survival
<i>Acer nigrum</i>	20			0.00
<i>Amelanchier arborea</i>	30	28	4	13.3
<i>Carya ovata</i>	20			0.0
<i>Celtis occidentalis</i>	15	7	7	46.7
<i>Cephalanthus occidentalis</i>	50	15	8	16.0
<i>Cornus amomum</i>	25			0.0
<i>Cornus stolonifera</i>	50	4	3	6.0
<i>Corylus americana</i>	50	11	14	28.0
<i>Fraxinus pennsylvanica</i>	25		17	68.0
<i>Juglans nigra</i>	20	10	16	80.0
<i>Physocarpus opulifolius</i>	25	7	1	4.0
<i>Platanus occidentalis</i>	10	4	1	10.0
<i>Quercus bicolor</i>	25	25	13	52.0
<i>Quercus macrocarpa</i>	25	16	18	72.0
<i>Salix amygdaloides</i>	30			0.0
<i>Salix nigra</i>	30			0.0
<i>Tilia Americana</i>	15	2	1	6.7
<i>Ulmus Americana</i>	5			0.0
Total	470	129	103	21.9

DISCUSSION

After three years of site development since construction, these sites showed limited progress towards wetland establishment. There is a high probability that Site 3 and a portion of Site 1 (Site 1A) will comply with project goal 1 (establishment of a jurisdictional wetland) by the end of the monitoring period. However, Site 1B and Site 2 lacked dominant hydrophytic vegetation, hydric soils and wetland hydrology, and are unlikely to develop these characteristics given current site conditions. Based on dominance of hydrophytic vegetation and presence of hydric soils, we estimate that 0.4 ha of 0.4 ha (1.0 ac of 1.0 ac) at Site 3 and 1.7 ha of 4.2 ha (4.2 ac of 10.4 ac) at Site 1 currently meet the jurisdictional criteria of a wetland (Fig. 2, Appendix B). The portion of Site 1 estimated to meet jurisdictional criteria based on vegetation and soils (i.e. Site 1A, Fig. 2, Appendix B) corresponded well to the estimated portion of the site meeting the wetland hydrology criterion (crosshatched area, Fig. 1, Appendix B).

Success was limited with respect to project goal 2 (minimum standards for floristic integrity and species composition). In both 2005 and 2006 the project area as a whole met performance standards for having an FQI above 20, when planted species are included, and for supporting greater than 30 native species. Floristic quality values generally increased from 2005 levels. However, for an area this large these performance standards should be considered low, and not necessarily indicative of successful restoration. Even highly disturbed or early successional sites of this size are likely to have more than 30 native species. All sites met the criterion of having a mean wetness coefficient less than zero, as well as the criterion of having greater than 60% vegetation cover. Only Site 3 met the criterion regarding dominance by non-native and weedy species. Because these sites were recently established they are dominated, in part, by disturbance-adapted, annual species (such as *Setaria faberi* and *Chenopodium album*). These species will decline in importance over the course of site development as they are replaced by perennial vegetation. However, the current dominance of Sites 1A and 1B by the aggressive, perennial, exotic grass *Phalaris arundinacea* will present a major barrier to establishing minimum standards for floristic integrity and species composition at the site. *Phalaris arundinacea* was also common at Site 3, and has increased in importance at that site since 2005. Encroachment by *P. arundinacea* may lead to a decrease in species diversity and FQI, and should be considered a threat to the success of these sites.

Several planted herbaceous species were located at each site, and planted species appeared to be spreading from the areas where they were originally planted. Continued persistence of planted herbaceous species will elevate the floristic integrity of these sites.

Site 2, the forested wetland, did not meet the planted tree and shrub survival criterion for project goal 3 in 2005 or 2006. Percent survival was far less than the goal of 80% survival, and several surviving trees and shrubs were overtopped by tall herbaceous vegetation. Because this site is not likely to meet the primary goal of creating jurisdictional wetland, it may not be practical to attempt to replant trees at this site.

Literature Cited

- Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. U.S. Army Engineer Waterways Experiment Station, Vicksburg, Mississippi. Technical Report Y-87-1.
- Federal Interagency Committee for Wetland Delineation. 1989. Federal manual for identifying and delineating jurisdictional wetlands. U.S. Army Corps of Engineers, U.S. Environmental Protection Agency, U.S. Fish and Wildlife Service, and U.S.D.A. Soil Conservation Service, Washington, D.C. Cooperative technical publication.
- Fucciolo, C. S., S. E. Benton, K. W. Carr, C. W. Knight, J. J. Miner, E. T. Plankell, G. E. Pociask, B. J. R. Sperling. 2006. Annual report for active IDOT wetland compensation and hydrologic monitoring sites. Report submitted to the Illinois Department of Transportation, Bureau of Design and Environment, Wetlands Unit.
- Grantham, D.R. 1980. Soil Survey of Winnebago and Boone Counties, Illinois. United States Department of Agriculture-Soil Conservation Service in cooperation with Illinois Agricultural Experiment Station. Illinois Agricultural Experiment Station Soil Report No. 107. 279 pp. + maps.
- Ogata, K.M. 1975. Drainage areas for Illinois streams. U.S. Geological Survey in cooperation with Illinois Institute for Environmental Quality. Water Resources Investigations 13-75. 120 pp.
- Plankell, E. T. and S. E. Benton. 2006. Pecatonica River Forest Preserve wetland compensation site. Report submitted to the Illinois Department of Transportation, Bureau of Design and Environment, Natural Resources Unit.
- Reed, P. B., Jr. 1988. National list of plant species that occur in wetlands: Illinois. U.S. Fish and Wildlife Service, National Wetlands Inventory. NERC-88/18.13.
- Swink, F., and G. Wilhelm. 1994. Plants of the Chicago region. Indiana Academy of Science, Indianapolis.
- Taft, J. B., G. S. Wilhelm, D. M. Ladd, and L. A. Masters. 1997. Floristic quality assessment for vegetation in Illinois - a method for assessing vegetation integrity. *Erigenia* 15:3-95.

APPENDIX A: WETLAND DETERMINATION FORMS

ROUTINE ON-SITE WETLAND DETERMINATION

Site 1A (page 2 of 5)

Field Investigators: Matthews, Kurylo, Zylka and Drahiem

Date: 14 August 2006

Project Name: Harrison Avenue Extension at Pecatonica

State: Illinois **County:** Winnebago **Applicant:** IDOT District 2

Site Name: Marsh/wet meadow

Legal Description: NE/4, NW/4, Sect. 19, T 27N, R 10E

Location: This site is located in northern part of the project area, just south of Blair Road.

HYDROLOGY

Inundated: Yes: X (partially) No: Depth of standing water: 0.3 m (1 ft)

Depth to saturated soil: from surface to 0.7 m (2.3 ft)

Overview of hydrological flow through the system: This site receives water through precipitation, sheet flow from surrounding higher ground, a culvert under Blair Road to the north, and possibly infrequent overflow from the Pecatonica River. Water leaves the site via evapotranspiration.

Size of Watershed: 4429 km² (1710 mi²) (Ogata 1975)

Other field evidence observed: None

Wetland hydrology: Yes: X No:

Rationale: This site is located in a depressional area. According to a report by ISGS personnel (Plankell and Benton 2006) this site was inundated or saturated for a sufficient duration to satisfy the wetland hydrology criterion during 2006.

DETERMINATION AND RATIONALE:

Is the site a wetland? Yes: X No:

Rationale: This site supports dominant hydrophytic vegetation, hydric soils, and wetland hydrology. We determined that this site is a wetland.

Determined by: Jeff Matthews and Jason Zylka (vegetation and hydrology)
Jessica Kurylo and Ian Drahiem (soils and hydrology)
Illinois Natural History Survey
1816 S. Oak Street
Champaign, Illinois 61820
(217) 244-2168 (Matthews)

ROUTINE ON-SITE WETLAND DETERMINATION

Site 1A (page 3 of 5)

Field Investigators: Matthews, Kurylo, Zylka and Drahiem

Date: 14 August 2006

Project Name: Harrison Avenue Extension at Pecatonica

State: Illinois **County:** Winnebago **Applicant:** IDOT District 2

Site Name: Marsh/wet meadow

Legal Description: NE/4, NW/4, Sect. 19, T 27N, R 10E

Location: This site is located in northern part of the project area, just south of Blair Road.

SPECIES LIST

Scientific Name	Common Name	Stratum	Wetland indicator status	C**
<i>Abutilon theophrasti</i>	velvet-leaf	herb	FACU-	*
<i>Acalypha rhomboidea</i>	three-seeded mercury	herb	FACU	0
<i>Acer saccharinum</i>	silver maple	tree	FACW	1
† <i>Acorus calamus</i>	sweetflag	herb	OBL	4
† <i>Alisma plantago-aquatica</i>	broad-leaf water-plantain	herb	OBL	2
<i>Amaranthus tuberculatus</i>	tall waterhemp	herb	OBL	1
<i>Ammannia coccinea</i>	long-leaved ammannia	herb	OBL	5
† <i>Andropogon gerardii</i>	big bluestem	herb	FAC-	5
<i>Asclepias incarnata</i>	swamp milkweed	herb	OBL	4
<i>Bidens cernua</i>	nodding beggar's ticks	herb	OBL	2
<i>Bidens frondosa</i>	common beggar's ticks	herb	FACW	1
<i>Carex comosa</i>	bristly sedge	herb	OBL	6
† <i>Carex hystricina</i>	bottlebrush sedge	herb	OBL	6
<i>Carex</i> sp.	sedge	herb	----	--
<i>Chamaesyce maculata</i>	nodding spurge	herb	FACU-	0
<i>Chenopodium album</i>	lamb's quarters	herb	FAC-	*
<i>Cirsium arvense</i>	Canada thistle	herb	FACU	*
<i>Cirsium vulgare</i>	bull thistle	herb	FACU-	*
<i>Cyperus ferruginescens</i>	rusty nut-sedge	herb	OBL	1
<i>Echinochloa muricata</i>	barnyard grass	herb	OBL	0
<i>Echinocystis lobata</i>	wild balsam-apple	herb	FACW-	4
<i>Epilobium coloratum</i>	cinnamon willow herb	herb	OBL	3
<i>Equisetum hyemale affine</i>	tall scouring rush	herb	FACW-	2
<i>Eragrostis pectinacea</i>	Carolina love grass	herb	FAC	0
<i>Erechtites hieracifolia</i>	fire weed	herb	FACU	2
<i>Erigeron annuus</i>	annual fleabane	herb	FAC-	1
<i>Eupatorium perfoliatum</i>	common boneset	herb	FACW+	4
<i>Helianthus tuberosus</i>	Jerusalem artichoke	herb	FAC	3
<i>Impatiens capensis</i>	jewelweed	herb	FACW	2
<i>Ipomoea pandurata</i>	wild sweet potato vine	herb	FACU	2
† <i>Iris shrevei</i>	southern blue flag	herb	OBL	5
<i>Leersia oryzoides</i>	rice cutgrass	herb	OBL	3
<i>Lemna minor</i>	common duckweed	herb	OBL	3

(continued on next page)

ROUTINE ON-SITE WETLAND DETERMINATION

Site 1A (page 4 of 5)

Field Investigators: Matthews, Kurylo, Zylka and Drahiem

Date: 14 August 2006

Project Name: Harrison Avenue Extension at Pecatonica

State: Illinois **County:** Winnebago **Applicant:** IDOT District 2

Site Name: Marsh/wet meadow

Legal Description: NE/4, NW/4, Sect. 19, T 27N, R 10E

Location: This site is located in northern part of the project area, just south of Blair Road.

SPECIES LIST (continued)

Scientific Name	Common Name	Stratum	Wetland indicator status	C**
<i>Lindernia dubia</i>	false pimpernel	herb	OBL	5
<i>Lythrum salicaria</i>	purple loosestrife	herb	OBL	*
† <i>Mimulus ringens</i>	monkey flower	herb	OBL	5
<i>Nasturtium officinale</i>	water cress	herb	OBL	*
<i>Oenothera biennis</i>	evening primrose	herb	FACU	1
<i>Panicum capillare</i>	witch grass	herb	FAC	0
<i>Panicum dichotomiflorum</i>	fall panicum	herb	FACW-	0
<i>Penthorum sedoides</i>	ditch stonecrop	herb	OBL	2
<i>Phalaris arundinacea</i>	reed canary grass	herb	FACW+	*
<i>Polygonum amphibium</i>	water smartweed	herb	OBL	3
<i>Polygonum lapathifolium</i>	curttop lady's thumb	herb	FACW+	0
<i>Polygonum pensylvanicum</i>	giant smartweed	herb	FACW+	1
<i>Polygonum persicaria</i>	spotted lady's thumb	herb	FACW	*
<i>Populus deltoides</i>	eastern cottonwood	shrub	FAC+	2
<i>Potentilla norvegica</i>	rough cinquefoil	herb	FAC	0
<i>Rotala ramosior</i>	tooth-cup	herb	OBL	4
<i>Salix exigua</i>	sandbar willow	shrub	OBL	1
† <i>Scirpus acutus</i>	great bulrush	herb	OBL	6
† <i>Scirpus atrovirens</i>	dark green bulrush	herb	OBL	4
<i>Scirpus tabernaemontanii</i>	great bulrush	herb	OBL	4
<i>Setaria faberi</i>	giant foxtail	herb	FACU+	*
<i>Setaria glauca</i>	pigeon grass	herb	FAC	*
† <i>Scutellaria lateriflora</i>	mad-dog skullcap	herb	OBL	4
<i>Solidago canadensis</i>	Canada goldenrod	herb	FACU	1
† <i>Solidago gigantea</i>	late goldenrod	herb	FACW	3
<i>Sonchus asper</i>	prickly sowthistle	herb	FAC	*
<i>Toxicodendron radicans</i>	poison ivy	shrub	FAC+	1
<i>Tridens flavus</i>	common purple top	herb	UPL	1
<i>Typha angustifolia</i>	narrow-leaved cattail	herb	OBL	*
<i>Typha latifolia</i>	cattail	herb	OBL	1
<i>Urtica dioica</i>	stinging nettle	herb	FAC+	2
<i>Verbena hastata</i>	blue vervain	herb	FACW+	3
<i>Verbena urticifolia</i>	white vervain	herb	FAC+	3

(continued on next page)

ROUTINE ON-SITE WETLAND DETERMINATION

Site 1A (page 5 of 5)

Field Investigators: Matthews, Kurylo, Zylka and Drahiem

Date: 14 August 2006

Project Name: Harrison Avenue Extension at Pecatonica

State: Illinois **County:** Winnebago **Applicant:** IDOT District 2

Site Name: Marsh/wet meadow

Legal Description: NE/4, NW/4, Sect. 19, T 27N, R 10E

Location: This site is located in northern part of the project area, just south of Blair Road.

SPECIES LIST (continued)

Scientific Name	Common Name	Stratum	Wetland indicator status	C**
<i>Vitis riparia</i>	riverbank grape	herb	FACW-	2
<i>Xanthium strumarium</i>	cocklebur	herb	FAC	0

** Coefficient of Conservatism (Taft et al. 1997)

* Non-native species

† Planted species

With planted species:

$$mCv = \sum C/N = 131/55 = 2.4$$

$$FQI = \sum C/\sqrt{N} = 131/\sqrt{55} = 17.7$$

Without planted species:

$$mCv = \sum C/N = 87/45 = 1.9$$

$$FQI = \sum C/\sqrt{N} = 87/\sqrt{45} = 13.0$$

ROUTINE ON-SITE WETLAND DETERMINATION

Site 1B (page 3 of 5)

Field Investigators: Matthews, Kurylo, Zylka and Drahiem

Date: 14 August 2006

Project Name: Harrison Avenue Extension at Pecatonica

State: Illinois **County:** Winnebago **Applicant:** IDOT District 2

Site Name: Non-native grassland

Legal Description: NE/4, NW/4, Sect. 19, T 27N, R 10E

Location: This site is located in northern part of the project area, just south of Blair Road, and north of the Pecatonica River.

SPECIES LIST

Scientific Name	Common Name	Stratum	Wetland indicator status	C**
<i>Abutilon theophrasti</i>	velvet-leaf	herb	FACU-	*
<i>Acalypha rhomboidea</i>	three-seeded mercury	herb	FACU	0
<i>Acer negundo</i>	box elder	herb, shrub	FACW-	1
<i>Acer saccharinum</i>	silver maple	herb	FACW	1
<i>Amaranthus tuberculatus</i>	tall waterhemp	herb	OBL	1
<i>Ambrosia trifida</i>	giant ragweed	herb	FAC+	0
† <i>Andropogon gerardii</i>	big bluestem	herb	FAC-	5
† <i>Anemone canadensis</i>	meadow anemone	herb	FACW	4
<i>Apocynum cannabinum</i>	dogbane	herb	FAC	2
<i>Arctium minus</i>	common burdock	herb	UPL	*
<i>Asclepias incarnata</i>	swamp milkweed	herb	OBL	4
<i>Asclepias syriaca</i>	common milkweed	herb	UPL	0
<i>Aster ontarionis</i>	Ontario aster	herb	FAC	4
<i>Aster pilosus</i>	hairy aster	herb	FACU+	0
<i>Bidens frondosa</i>	common beggar's ticks	herb	FACW	1
<i>Bidens vulgata</i>	tall beggar's ticks	herb	FACW	0
<i>Calystegia sepium</i>	American bindweed	herb	FAC	1
<i>Campanula americana</i>	American bellflower	herb	FAC	4
<i>Chenopodium album</i>	lamb's quarters	herb	FAC-	*
<i>Cirsium arvense</i>	Canada thistle	herb	FACU	*
<i>Cirsium discolor</i>	pasture thistle	herb	UPL	3
<i>Cirsium vulgare</i>	bull thistle	herb	FACU-	*
<i>Conyza canadensis</i>	horseweed	herb	FAC-	0
<i>Echinochloa muricata</i>	barnyard grass	herb	OBL	0
<i>Elymus virginicus</i>	Virginia wild rye	herb	FACW-	4
<i>Eragrostis pectinacea</i>	Carolina love grass	herb	FAC	0
<i>Erigeron annuus</i>	annual fleabane	herb	FAC-	1
<i>Geum canadense</i>	white avens	herb	FAC	2
<i>Geum laciniatum</i>	rough avens	herb	FACW	2
<i>Hackelia virginiana</i>	stickseed	herb	FAC-	1
† <i>Iris shrevei</i>	southern blue flag	herb	OBL	5
<i>Lactuca serriola</i>	prickly lettuce	herb	FAC	*
<i>Laportea canadensis</i>	wood nettle	herb	FACW	2

(continued on next page)

ROUTINE ON-SITE WETLAND DETERMINATION

Site 1B (page 4 of 5)

Field Investigators: Matthews, Kurylo, Zylka and Drahiem

Date: 14 August 2006

Project Name: Harrison Avenue Extension at Pecatonica

State: Illinois **County:** Winnebago **Applicant:** IDOT District 2

Site Name: Non-native grassland

Legal Description: NE/4, NW/4, Sect. 19, T 27N, R 10E

Location: This site is located in northern part of the project area, just south of Blair Road, and north of the Pecatonica River.

SPECIES LIST (continued)

Scientific Name	Common Name	Stratum	Wetland indicator status	C**
<i>Morus alba</i>	white mulberry	herb	FAC	*
<i>Oenothera biennis</i>	evening primrose	herb	FACU	1
<i>Oxalis stricta</i>	yellow wood sorrel	herb	FACU	0
<i>Panicum capillare</i>	witch grass	herb	FAC	0
<i>Panicum dichotomiflorum</i>	fall panicum	herb	FACW-	0
† <i>Panicum virgatum</i>	prairie switchgrass	herb	FAC+	4
<i>Phalaris arundinacea</i>	reed canary grass	herb	FACW+	*
<i>Physalis subglabrata</i>	smooth ground cherry	herb	UPL	0
<i>Pilea pumila</i>	Canada clearweed	herb	FACW	3
<i>Polygonum amphibium</i>	water smartweed	herb	OBL	3
<i>Polygonum pensylvanicum</i>	giant smartweed	herb	FACW+	1
<i>Polygonum persicaria</i>	spotted lady's thumb	herb	FACW	*
<i>Polygonum scandens</i>	climbing buckwheat	herb	FAC	2
<i>Populus deltoides</i>	eastern cottonwood	herb	FAC+	2
<i>Potentilla norvegica</i>	rough cinquefoil	herb	FAC	0
<i>Ratibida pinnata</i>	drooping coneflower	herb	UPL	4
<i>Rubus occidentalis</i>	black raspberry	shrub	UPL	2
<i>Rudbeckia hirta</i>	black-eyed Susan	herb	FACU	2
<i>Rumex crispus</i>	curly dock	herb	FAC+	*
<i>Sambucus canadensis</i>	common elder	shrub	FACW-	2
<i>Setaria faberi</i>	giant foxtail	herb	FACU+	*
<i>Setaria glauca</i>	pigeon grass	herb	FAC	*
† <i>Silphium perfoliatum</i>	cup plant	herb	FACW-	4
<i>Solanum carolinense</i>	horse nettle	herb	FACU-	0
<i>Solidago canadensis</i>	Canada goldenrod	herb	FACU	1
† <i>Solidago gigantea</i>	late goldenrod	herb	FACW	3
<i>Sonchus asper</i>	prickly sowthistle	herb	FAC	*
† <i>Spartina pectinata</i>	freshwater cord grass	herb	FACW+	4
<i>Stachys tenuifolia</i>	slenderleaf betony	herb	OBL	5
<i>Stellaria media</i>	common chickweed	herb	FACU	*
<i>Teucrium canadense</i>	American germander	herb	FACW-	3
<i>Toxicodendron radicans</i>	poison ivy	herb	FAC+	1
<i>Tridens flavus</i>	common purple top	herb	UPL	1

(continued on next page)

ROUTINE ON-SITE WETLAND DETERMINATION
Site 1B (page 5 of 5)

Field Investigators: Matthews, Kurylo, Zylka and Drahiem
Date: 14 August 2006
Project Name: Harrison Avenue Extension at Pecatonica
State: Illinois **County:** Winnebago **Applicant:** IDOT District 2
Site Name: Non-native grassland
Legal Description: NE/4, NW/4, Sect. 19, T 27N, R 10E
Location: This site is located in northern part of the project area, just south of Blair Road, and north of the Pecatonica River.

SPECIES LIST (continued)

Scientific Name	Common Name	Stratum	Wetland indicator status	C**
<i>Ulmus americana</i>	American elm	herb	FACW-	5
<i>Urtica dioica</i>	stinging nettle	herb	FAC+	2
<i>Veronica peregrina</i>	purslane speedwell	herb	FACW+	0
<i>Viola pratincola</i>	common blue violet	herb	FAC	1
<i>Vitis riparia</i>	riverbank grape	herb	FACW-	2

** Coefficient of Conservatism (Taft et al. 1997)

* Non-native species

† Planted species

With planted species:

$$mCv = \sum C/N = 106/57 = 1.9$$

$$FQI = \sum C/\sqrt{N} = 106/\sqrt{57} = 14.1$$

Without planted species:

$$mCv = \sum C/N = 77/50 = 1.5$$

$$FQI = \sum C/\sqrt{N} = 77/\sqrt{50} = 10.9$$

ROUTINE ON-SITE WETLAND DETERMINATION

Site 2 (page 3 of 5)

Field Investigators: Matthews, Kurylo, Zylka and Drahiem

Date: 14 August 2006

Project Name: Harrison Avenue Extension at Pecatonica

State: Illinois **County:** Winnebago **Applicant:** IDOT District 2

Site Name: Forested site

Legal Description: S/2, NW/4, Sect. 19, T 27N, R 10E

Location: This site is located in southern part of the project area, north and west of the Pecatonica River.

SPECIES LIST

Scientific Name	Common Name	Stratum	Wetland indicator status	C**
<i>Abutilon theophrasti</i>	velvet-leaf	herb	FACU-	*
<i>Acer negundo</i>	box elder	shrub	FACW-	1
<i>Acer saccharinum</i>	silver maple	herb	FACW	1
<i>Agropyron repens</i>	quack grass	herb	FACU	*
<i>Ambrosia artemisiifolia</i>	common ragweed	herb	FACU	0
<i>Ambrosia trifida</i>	giant ragweed	herb	FAC+	0
† <i>Amelanchier arborea</i>	juneberry	tree	FACU	7
† <i>Andropogon gerardii</i>	big bluestem	herb	FAC-	5
† <i>Angelica atropurpurea</i>	angelica	herb	OBL	6
<i>Apocynum cannabinum</i>	dogbane	herb	FAC	2
<i>Asclepias syriaca</i>	common milkweed	herb	UPL	0
<i>Aster ontarionis</i>	Ontario aster	herb	FAC	4
<i>Aster pilosus</i>	hairy aster	herb	FACU+	0
<i>Bidens vulgata</i>	tall beggar's ticks	herb	FACW	0
<i>Campanula americana</i>	American bellflower	herb	FAC	4
<i>Carex</i> sp.	sedge	herb	----	--
† <i>Carex vulpinoidea</i>	fox sedge	herb	OBL	3
† <i>Celtis occidentalis</i>	hackberry	tree	FAC-	3
† <i>Cephalanthus occidentalis</i>	buttonbush	shrub	OBL	4
<i>Chenopodium album</i>	lamb's quarters	herb	FAC-	*
<i>Cirsium arvense</i>	Canada thistle	herb	FACU	*
<i>Cirsium vulgare</i>	bull thistle	herb	FACU-	*
<i>Conyza canadensis</i>	horseweed	herb	FAC-	0
† <i>Cornus stolonifera</i>	red osier dogwood	shrub	FACW	4
† <i>Corylus americana</i>	American filbert	shrub	FACU-	4
<i>Elymus canadensis</i>	Canada wild rye	herb	FAC-	4
<i>Elymus virginicus</i>	Virginia wild rye	herb	FACW-	4
<i>Fraxinus pennsylvanica</i>	green ash	shrub, herb	FACW	2
<i>Geum canadense</i>	white avens	herb	FAC	2
<i>Hackelia virginiana</i>	stickseed	herb	FAC-	1
<i>Hordeum jubatum</i>	squirrel-tail	herb	FAC+	*
† <i>Iris shrevei</i>	southern blue flag	herb	OBL	5

(continued on next page)

ROUTINE ON-SITE WETLAND DETERMINATION

Site 2 (page 3 of 5)

Field Investigators: Matthews, Kurylo, Zylka and Drahiem

Date: 14 August 2006

Project Name: Harrison Avenue Extension at Pecatonica

State: Illinois **County:** Winnebago **Applicant:** IDOT District 2

Site Name: Forested site

Legal Description: S/2, NW/4, Sect. 19, T 27N, R 10E

Location: This site is located in southern part of the project area, north and west of the Pecatonica River.

SPECIES LIST (continued)

Scientific Name	Common Name	Stratum	Wetland indicator status	C**
† <i>Juglans nigra</i>	black walnut	tree	FACU	4
<i>Lactuca serriola</i>	prickly lettuce	herb	FAC	*
<i>Laportea canadensis</i>	wood nettle	herb	FACW	2
<i>Morus alba</i>	white mulberry	shrub, herb	FAC	*
<i>Oenothera biennis</i>	evening primrose	herb	FACU	1
<i>Oxalis stricta</i>	yellow wood sorrel	herb	FACU	0
† <i>Panicum virgatum</i>	prairie switchgrass	herb	FAC+	4
<i>Phalaris arundinacea</i>	reed canary grass	herb	FACW+	*
<i>Physalis subglabrata</i>	smooth ground cherry	herb	UPL	0
† <i>Physocarpus opulifolius</i>	common ninebark	shrub	FACW-	7
† <i>Platanus occidentalis</i>	sycamore	tree	FACW	3
<i>Polygonum amphibium</i>	water smartweed	herb	OBL	3
<i>Polygonum lapathifolium</i>	curttop lady's thumb	herb	FACW+	0
<i>Polygonum pensylvanicum</i>	giant smartweed	herb	FACW+	1
<i>Polygonum persicaria</i>	spotted lady's thumb	herb	FACW	*
<i>Polygonum scandens</i>	climbing buckwheat	herb	FAC	2
<i>Populus deltoides</i>	eastern cottonwood	herb	FAC+	2
<i>Potentilla norvegica</i>	rough cinquefoil	herb	FAC	0
† <i>Quercus bicolor</i>	swamp white oak	tree	FACW+	7
† <i>Quercus macrocarpa</i>	bur oak	tree	FAC-	5
<i>Rudbeckia laciniata</i>	cutleaf coneflower	herb	FACW+	3
<i>Rumex altissimus</i>	pale dock	herb	FACW-	2
<i>Rumex crispus</i>	curly dock	herb	FAC+	*
<i>Salix exigua</i>	sandbar willow	shrub	OBL	1
<i>Scrophularia marilandica</i>	late figwort	herb	FACU-	4
<i>Setaria faberi</i>	giant foxtail	herb	FACU+	*
† <i>Silphium perfoliatum</i>	cup plant	herb	FACW-	4
<i>Smilax hispida</i>	bristly greenbrier	herb	FAC	3
<i>Solanum carolinense</i>	horse nettle	herb	FACU-	0
<i>Solidago canadensis</i>	Canada goldenrod	herb	FACU	1
<i>Solidago gigantea</i>	late goldenrod	herb	FACW	3
<i>Sonchus asper</i>	prickly sowthistle	herb	FAC	*
<i>Taraxacum officinale</i>	common dandelion	herb	FACU	*

(continued on next page)

ROUTINE ON-SITE WETLAND DETERMINATION

Site 2 (page 5 of 5)

Field Investigators: Matthews, Kurylo, Zylka and Drahiem

Date: 14 August 2006

Project Name: Harrison Avenue Extension at Pecatonica

State: Illinois **County:** Winnebago **Applicant:** IDOT District 2

Site Name: Forested site

Legal Description: S/2, NW/4, Sect. 19, T 27N, R 10E

Location: This site is located in southern part of the project area, north and west of the Pecatonica River.

SPECIES LIST (continued)

Scientific Name	Common Name	Stratum	Wetland indicator status	C**
† <i>Tilia americana</i>	basswood	shrub	FACU	5
<i>Toxicodendron radicans</i>	poison ivy	herb	FAC+	1
<i>Tridens flavus</i>	common purple top	herb	UPL	1
<i>Ulmus americana</i>	American elm	herb	FACW-	5
<i>Verbena urticifolia</i>	white vervain	herb	FAC+	3
<i>Viola sororia</i>	woolly blue violet	herb	FAC-	3
<i>Vitis riparia</i>	riverbank grape	herb	FACW-	2

** Coefficient of Conservatism (Taft et al. 1997)

* Non-native species

† Planted species

With planted species:

$$mCv = \sum C/N = 148/57 = 2.6$$

$$FQI = \sum C/\sqrt{N} = 148/\sqrt{57} = 19.6$$

Without planted species:

$$mCv = \sum C/N = 68/40 = 1.7$$

$$FQI = \sum C/\sqrt{N} = 68/\sqrt{40} = 10.8$$

ROUTINE ON-SITE WETLAND DETERMINATION
Site 3 (page 2 of 4)

Field Investigators: Matthews, Kurylo, Zylka and Drahiem
Date: 14 August 2006
Project Name: Harrison Avenue Extension at Pecatonica
State: Illinois **County:** Winnebago **Applicant:** IDOT District 2
Site Name: Streambed wetland
Legal Description: NW/4, Sect. 19, T 27N, R 10E
Location: This site is located along the western boundary of the project area, beginning just south of Blair Road, and extending south to the Pecatonica River.

HYDROLOGY

Inundated: Yes: X No: Depth of standing water: 0.3 m (1 ft)

Depth to saturated soil: saturated at surface

Overview of hydrological flow through the system: This site is a streambed and receives water primarily through stream flow from the north. In addition the site receives water through precipitation, sheet flow from surrounding higher ground, and possible backflow from the Pecatonica River. Water leaves the site via evapotranspiration, soil infiltration and stream flow into the Pecatonica River at the south end.

Size of Watershed: 4429 km² (1710 mi²) (Ogata 1975)

Other field evidence observed: Barren, cracked soil in some areas

Wetland hydrology: Yes: X No:

Rationale: This site is located in a streambed and holds water for a long duration during the growing season. Although this site has not been monitored by the ISGS, field evidence indicates that the site was inundated or saturated for a sufficient duration to satisfy the wetland hydrology criterion during 2006.

DETERMINATION AND RATIONALE:

Is the site a wetland? Yes: X No:

Rationale: This site supports dominant hydrophytic vegetation, hydric soils, and wetland hydrology. We determined that this site is a wetland.

Determined by: Jeff Matthews and Jason Zylka (vegetation and hydrology)
Jessica Kurylo and Ian Drahiem (soils and hydrology)
Illinois Natural History Survey
1816 S. Oak Street
Champaign, Illinois 61820
(217) 244-2168 (Matthews)

ROUTINE ON-SITE WETLAND DETERMINATION

Site 3 (page 3 of 4)

Field Investigators: Matthews, Kurylo, Zylka and Drahiem

Date: 14 August 2006

Project Name: Harrison Avenue Extension at Pecatonica

State: Illinois **County:** Winnebago **Applicant:** IDOT District 2

Site Name: Streambed wetland

Legal Description: NW/4, Sect. 19, T 27N, R 10E

Location: This site is located along the western boundary of the project area, beginning just south of Blair Road, and extending south to the Pecatonica River.

SPECIES LIST

Scientific Name	Common Name	Stratum	Wetland indicator status	C**
<i>Abutilon theophrasti</i>	velvet-leaf	herb	FACU-	*
<i>Acer saccharinum</i>	silver maple	tree, herb	FACW	1
<i>Alisma plantago-aquatica</i>	broad-leaf water-plantain	herb	OBL	2
<i>Amaranthus tuberculatus</i>	tall waterhemp	herb	OBL	1
<i>Bidens cernua</i>	nodding beggar's ticks	herb	OBL	2
<i>Bidens frondosa</i>	common beggar's ticks	herb	FACW	1
<i>Cephalanthus occidentalis</i>	buttonbush	shrub	OBL	4
<i>Echinochloa muricata</i>	barnyard grass	herb	OBL	0
<i>Eleocharis erythropoda</i>	spike rush	herb	OBL	3
<i>Elymus virginicus</i>	Virginia wild rye	herb	FACW-	4
<i>Impatiens capensis</i>	jewelweed	herb	FACW	2
<i>Leersia oryzoides</i>	rice cutgrass	herb	OBL	3
<i>Lemna minor</i>	common duckweed	herb	OBL	3
<i>Panicum capillare</i>	witch grass	herb	FAC	0
<i>Phalaris arundinacea</i>	reed canary grass	herb	FACW+	*
<i>Pilea pumila</i>	Canada clearweed	herb	FACW	3
<i>Polygonum amphibium</i>	water smartweed	herb	OBL	3
<i>Polygonum lapathifolium</i>	curttop lady's thumb	herb	FACW+	0
<i>Polygonum pensylvanicum</i>	giant smartweed	herb	FACW+	1
<i>Rumex crispus</i>	curly dock	herb	FAC+	*
† <i>Sagittaria latifolia</i>	arrowhead	herb	OBL	4
<i>Salix exigua</i>	sandbar willow	shrub	OBL	1
<i>Salix nigra</i>	black willow	tree	OBL	3
† <i>Scirpus acutus</i>	great bulrush	herb	OBL	6
<i>Scirpus atrovirens</i>	dark green bulrush	herb	OBL	4
<i>Scirpus cyperinus</i>	wool grass	herb	OBL	5
† <i>Scirpus fluviatilis</i>	river bulrush	herb	OBL	3
<i>Scirpus tabernaemontanii</i>	great bulrush	herb	OBL	4
<i>Scutellaria lateriflora</i>	mad-dog skullcap	herb	OBL	4
<i>Setaria faberi</i>	giant foxtail	herb	FACU+	*
† <i>Sparganium eurycarpum</i>	burreed	herb	OBL	5
<i>Stachys tenuifolia</i>	slenderleaf betony	herb	OBL	5

(continued on next page)

ROUTINE ON-SITE WETLAND DETERMINATION

Site 3 (page 4 of 4)

Field Investigators: Matthews, Kurylo, Zylka and Drahiem

Date: 14 August 2006

Project Name: Harrison Avenue Extension at Pecatonica

State: Illinois **County:** Winnebago **Applicant:** IDOT District 2

Site Name: Streambed wetland

Legal Description: NW/4, Sect. 19, T 27N, R 10E

Location: This site is located along the western boundary of the project area, beginning just south of Blair Road, and extending south to the Pecatonica River.

SPECIES LIST (continued)

Scientific Name	Common Name	Stratum	Wetland indicator status	C**
<i>Ulmus americana</i>	American elm	tree	FACW-	5
<i>Urtica dioica</i>	stinging nettle	herb	FAC+	2
<i>Verbena hastata</i>	blue vervain	herb	FACW+	3
<i>Verbena urticifolia</i>	white vervain	herb	FAC+	3
<i>Vitis riparia</i>	riverbank grape	vine	FACW-	2
<i>Xanthium strumarium</i>	cocklebur	herb	FAC	0

** Coefficient of Conservatism (Taft et al. 1997)

* Non-native species

† Planted species

With planted species:

$$mCv = \sum C/N = 92/34 = 2.7$$

$$FQI = \sum C/\sqrt{N} = 92/\sqrt{34} = 15.8$$

Without planted species:

$$mCv = \sum C/N = 74/30 = 2.5$$

$$FQI = \sum C/\sqrt{N} = 74/\sqrt{30} = 13.5$$

APPENDIX B: VEGETATION, WETLAND AND HYDROLOGIC MAPS

Figure 1: Estimated extent of 2006 wetland hydrology within the project area (figure prepared by ISGS, from Plankell and Benton 2006).





Figure 2: Locations of wetland determination sites/vegetation communities within the project area.



APPENDIX C: RESULTS OF QUANTITATIVE VEGETATION SAMPLING

Table 1: Results of 2006 quantitative vegetation sampling at Site 1A (marsh/wet meadow)

Species	Frequency	Relative frequency	Average cover	Relative cover	IV
<i>Phalaris arundinacea</i>	100.0	36.7	69.6	67.8	52.3
<i>Polygonum amphibium</i>	54.5	20.0	12.4	12.1	16.0
<i>Echinochloa muricata</i>	9.1	3.3	7.7	7.5	5.4
<i>Leersia oryzoides</i>	9.1	3.3	5.7	5.5	4.4
<i>Setaria faberi</i>	18.2	6.7	1.6	1.6	4.1
<i>Bidens cernua</i>	9.1	3.3	3.4	3.3	3.3
<i>Abutilon theophrasti</i>	9.1	3.3	0.3	0.3	1.8
<i>Chenopodium album</i>	9.1	3.3	0.3	0.3	1.8
<i>Erigeron annuus</i>	9.1	3.3	0.3	0.3	1.8
<i>Polygonum pennsylvanicum</i>	9.1	3.3	0.3	0.3	1.8
<i>Potentilla norvegica</i>	9.1	3.3	0.3	0.3	1.8
<i>Scutellaria lateriflora</i>	9.1	3.3	0.3	0.3	1.8
<i>Rotala ramosior</i>	9.1	3.3	0.3	0.3	1.8
<i>Vitis riparia</i>	9.1	3.3	0.3	0.3	1.8
Sum	273	100	103	100	100

Table 2: Results of 2006 quantitative vegetation sampling at Site 1B (non-native grassland)

Species	Frequency	Relative frequency	Average cover	Relative cover	IV
<i>Setaria faberi</i>	92.3	17.9	43.5	49.3	33.6
<i>Phalaris arundinacea</i>	30.8	6.0	9.8	11.1	8.5
<i>Chenopodium album</i>	61.5	11.9	2.8	3.1	7.5
<i>Polygonum pennsylvanicum</i>	30.8	6.0	6.3	7.2	6.6
<i>Lactuca serriola</i>	38.5	7.5	3.9	4.4	6.0
<i>Abutilon theophrasti</i>	53.8	10.4	1.2	1.3	5.9
<i>Echinochloa muricata</i>	23.1	4.5	5.3	6.0	5.2
<i>Setaria glauca</i>	15.4	3.0	4.0	4.6	3.8
<i>Veronica perigrina</i>	30.8	6.0	0.9	1.0	3.5
<i>Conyza canadensis</i>	23.1	4.5	1.6	1.8	3.2
<i>Potentilla norvegica</i>	23.1	4.5	0.7	0.8	2.6
<i>Solidago canadensis</i>	7.7	1.5	2.9	3.3	2.4
<i>Panicum capillare</i>	7.7	1.5	1.2	1.3	1.4
<i>Solanum carolinense</i>	7.7	1.5	1.2	1.3	1.4
<i>Ulmus americana</i>	7.7	1.5	1.2	1.3	1.4
<i>Acer saccharinum</i>	7.7	1.5	0.2	0.3	0.9
<i>Amaranthus tuberculatus</i>	7.7	1.5	0.2	0.3	0.9
<i>Bidens vulgata</i>	7.7	1.5	0.2	0.3	0.9
<i>Cirsium arvense</i>	7.7	1.5	0.2	0.3	0.9
<i>Erigeron annuus</i>	7.7	1.5	0.2	0.3	0.9
<i>Polygonum persicaria</i>	7.7	1.5	0.2	0.3	0.9
<i>Rumex crispus</i>	7.7	1.5	0.2	0.3	0.9
<i>Spartina pectinata</i>	7.7	1.5	0.2	0.3	0.9
Sum	515	100	88	100	100

Table 3: Results of 2006 vegetation sampling at Site 3 (streambed wetland)

Species	Commonness
<i>Bidens frondosa</i>	5
<i>Leersia oryzoides</i>	5
<i>Lemna minor</i>	5
<i>Bidens cernua</i>	4
<i>Phalaris arundinacea</i>	4
<i>Sagittaria latifolia</i>	4
<i>Acer saccharinum</i>	3
<i>Echinochloa muricata</i>	3
<i>Eleocharis erythropoda</i>	3
<i>Alisma plantago-aquatica</i>	2
<i>Impatiens capensis</i>	2
<i>Polygonum pennsylvanicum</i>	2
<i>Salix exigua</i>	2
<i>Scirpus cyperinus</i>	2
<i>Scirpus fluviatilis</i>	2
<i>Scirpus tabernaemontanii</i>	2
<i>Sparganium eurycarpum</i>	2
<i>Stachys tenuifolia</i>	2
<i>Verbena hastata</i>	2
<i>Xanthium strumarium</i>	2
<i>Abutilon theophrasti</i>	1
<i>Amaranthus tuberculatus</i>	1
<i>Cephalanthus occidentalis</i>	1
<i>Elymus virginicus</i>	1
<i>Panicum capillare</i>	1
<i>Pilea pumila</i>	1
<i>Polygonum amphibium</i>	1
<i>Polygonum lapathifolium</i>	1
<i>Rumex crispus</i>	1
<i>Salix nigra</i>	1
<i>Scirpus acutus</i>	1
<i>Scirpus atrovirens</i>	1
<i>Scutellaria lateriflora</i>	1
<i>Setaria faberi</i>	1
<i>Ulmus americana</i>	1
<i>Urtica dioica</i>	1
<i>Verbena urticifolia</i>	1
<i>Vitis riparia</i>	1

APPENDIX D: PHOTOGRAPHS OF WETLAND MITIGATION SITES

Fig. 1. View of Sites 1A and 1B from northeast corner of 1A, facing south



Fig. 2. View of Site 1A from northwest corner, facing southeast



Fig. 3. View of the boundary between Site 1A (left) and Site 1B (right), facing northeast



Fig. 4. View of Site 1B from southwest corner, facing east



Fig. 5. View of Site 2 from northwest corner, facing southeast



Fig. 6. View of Site 2 from southeast corner, facing northwest



Fig. 7. View of Site 3 from center, facing north



Fig. 8. View of Site 3 from center, facing south

